

Glia-Neuron GPCR Signaling as a Therapeutic Route to Axon Regeneration

cAMP-PKA Activation Elevates Excitability and Regrowth in Neurons

VISUAL ABSTRACT




Figure 1: Visual Abstract. Image created by author using Biorender, 2025

Background




Figure 3: PGE₂ Signaling Schematic. Image created by author using Biorender, 2025

INTRODUCTION

The Problem




Figure 2: Problem Statement. Image created by author using Biorender and Canva, 2025

METHODS




Figure 4: Overview of Methodology. Image created by author using Biorender and Canva, 2025

Research Questions

- Can PGE₂ promote axonal re-extension and cell survival after injury?
- What is the optimal dosage of PGE₂ to prevent toxicity and maximize protective effects?
- How does PGE₂ impact neuronal excitability after injury?
- How do PGE₂ downstream signaling molecules and EP Receptors affect the efficacy of PGE₂ in promoting a repair state?

RESULTS

dmPGE₂ Promoted Re-extension





Figure 5A-B: dmPGE₂ boosts re-extension. Image created by author using Prism, 2025

126.29% more growth in dmPGE₂ wells compared to DMSO control


Optimal Dosage of PGE₂ Improves Survival After Injury

Dead/Alive Cells: 24hr After Injury



1µM and 10µM dosages significantly improved survival

Molecular Docking



Binding Affinity
-8.28 kcal/mol

Table 1: Binding Affinity. Created by author using Canva, 2025

LAT1 + prodrug shows potential for delivery

Figure 7: Prodrug docked to LAT1. Image created by author using ChimeraX, 2025

Brainbow AAV: More Axonal Regrowth Following Axotomy




Figure 8: Brainbow sample images. Image created by author using Fiji, 2025

PGE₂/cAMP/MAPK Differences Across Injury and Naive




Figure 10: DEG and Pathway Analysis Results. Image created by author using R (GGPlots) and Appyter, 2025

cAMP downregulated in apoptotic neurons, MAPK and cAMP upregulated in Schwann cells after injury

Increased Voltage-Gated Sodium Channel Activation in dmPGE₂-treated Cells




Figure 9: RNAScope Quantification Results. Image created by author using Prism, 2025

IMPACT

- Offers direction for future treatment - PGE₂ may be a useful **therapeutic target** to promote the re-extension of axons following injury
- Axonal re-extension can not only **prevent cell death**, but also lead to a **restoration of function** for millions of patients

KEY REFERENCES

- Aid, S., Langenbach, R., & Bossert, F. (2000). Neuroinflammatory response to lipopolysaccharide is exacerbated in mice genetically deficient in cyclooxygenase-2. *Journal of Neuroinflammation*, 1(1). <https://doi.org/10.1186/1475-2875-1-1>
- Feng, Z., & Gao, F. (2012). Stem cell challenges in the treatment of neurodegenerative disease. *Cell neuroscience & therapeutics*, 1(2), 142-148. <https://doi.org/10.1155/1530-8849/2011/2/142>
- Hultman, J., Pehkonen, S., Ojanne, M., Hultman, T., Hultman, M., Aarås, S., Raponen, M., Hakonen, K. S., & Hultman, K. M. (2019). 1-Type Amino Acid Transporter 1 (LAT1) is a Novel Prodrug Carrier for the Delivery of Drugs into Neurons. *Neuroscience Letters*, 691, 1-5. <https://doi.org/10.1016/j.neulet.2019.04.009>
- Katani, H., Obara, P. S., Thirumangalakudi, A. P., O'Connell, E. M., Iyer, M., Donovan, L. J., Dugan, M. Q., Ambal, N., Granato, A., Zeng, H., Saw, N. L., Brossid, L., A., Sloan, S. A., Gray, E. E., Tran, K. V., Vishni, A., Yeh, A. K., Manich, A. E., Huber, M., Agrawal, A., ... Zuhars, J. B. (2024). Schwann cell-associated PGE₂ promotes sensory neuron excitability during development. *Cell*, 127(1), 460-472. <https://doi.org/10.1016/j.cell.2024.07.023>
- Mohr, S., Gonen, C., Ahmed, I. Z., Wang, M., & Hultman, K. M. (2023). Neurodegenerative disorders: Assessing the impact of natural vs drug-induced treatment options. *Agry medicine (Milton N.S.W.)*, 6(1), 62-67. <https://doi.org/10.1002/agry.12343>
- Morales, A. B., Järvinen, J., Löffler, S., Hultman, J., Aarås, S., Lehtonen, M., Jalkanen, A., & Hultman, K. M. (2020). L-Type Amino Acid Transporter 1 Enables the Efficient Brain Delivery of Small-Molecule Prodrugs across the Blood-Brain Barrier and into Human and Mouse Brain Parenchymal Cells. *ACS chemical neuroscience*, 11(24), 4301-4313. <https://doi.org/10.1021/acschemneuro.0c00564>
- Wang, K., Wang, S., Chen, Y., Wu, D., Hu, X., Lu, Y., Wang, L., Bao, L., Li, C., & Zhang, X. (2021). Single-cell transcriptomic analysis of somatosensory neurons uncovers temporal development of neuropathic pain. *Cell research*, 31(8), 904-918. <https://doi.org/10.1038/s41422-021-00479-9>
- Yokoyama, U., Inaba, K., Umemura, M., Fujita, T., & Ishikawa, Y. (2013). The prostaglandin EP4 receptor and its signaling pathway. *Pharmacological reviews*, 65(3), 1010-1032. <https://doi.org/10.1124/pr.112.001916>